

# FEED RATION GUIDELINES



## Aquatic and Crustaceans

Generally, the addition of AZOMITE® at 0.5% is suggested. Shrimp and tilapia studies using from 0.3% to 1.0% have all reported significant increase in weight gain.

## Beef

We recommend adding AZOMITE® at 1/2% (10 pounds per ton) of the total prepared feed mix. A maximum of 2% (40 pounds per ton) may be added in accordance with good feed manufacturing or feeding practices or offered “free choice”.

## Dairy Cows

We recommend adding 2-4 ounces of AZOMITE® to the daily rations of dairy cows.

## Horses

We recommend adding 2-4 ounces of AZOMITE® to the daily rations of horses.

## Poultry

For poultry on a prepared feed diet, we recommend adding AZOMITE® at 1/2% (10 pounds per ton) of the total feed mix. A maximum of 2% (40 pounds per ton) may be added in accordance with good feed manufacturing or feeding practices.

## Sheep

For livestock on a prepared feed diet, we generally recommend adding AZOMITE® at 1/2% (10 pounds per ton) of the total feed mix. A maximum of 2% (40 pounds per ton) may be added in accordance with good feed manufacturing or feeding practices.

## Swine

For livestock on a prepared feed diet, we generally recommend adding AZOMITE® at 1/2% (10 pounds per ton) of the total feed mix. A maximum of 2% (40 pounds per ton) may be added in accordance with good feed manufacturing or feeding practices.

